

**Neuropathic pain** is commonly described as **numbness, tingling, burning, stabbing, shooting.**

- Patients with CKD may experience pain from a variety of causes. They might have neuropathic and/or nociceptive (musculoskeletal) pain.
- Consider using the Follow-up Pain Assessment Tool weekly to monitor effect of pain management.
- See: [Pain Patient Handout](#)

## ► Step 1: Adjuvant

### 1<sup>st</sup> Line:

Gabapentin 50-100\* mg PO nightly. If not effective, it can be further titrated by 100 mg every 7 nights to a maximum of 300 mg PO nightly. It should be taken 2-3 hours before bed due to delay of peak onset. *Note that gabapentin is not commercially available in 50 mg capsules, but can be compounded for patients if the recommended low starting dose is desired.* The most common side effects are drowsiness, dizziness, confusion, and fatigue. Peripheral edema may also be a side effect.

**or**

Pregabalin 25 mg PO nightly. Titrate by 25 mg every 7 nights to a maximum of 75 mg PO nightly. It should be taken 2 hours before bedtime. Similar to gabapentin, but more expensive and not covered by Seniors' or Basic Alberta Blue Cross plans. Other private plans may cover the cost. Potential side effects are similar to those of gabapentin.

### 2<sup>nd</sup> Line

TCA antidepressant (unless contraindications i.e. conduction abnormalities on ECG, or excess weight gain)

Amitriptyline (Elavil) 10-25 mg PO daily (max dose 75 mg daily). Titrate by 10-25 mg every week as required.

### Is pain now adequately controlled?

Yes: Reassess at least monthly using the Follow-Up Pain Assessment Tool

No benefit: STOP Adjuvant and START Non-Opioid/Weak Opioid (Step 2)

Some benefit but inadequate: ADD Non-Opioid/Weak Opioid (Step 2)

## ► Step 2: Non-Opioid

Due to the pharmacokinetics and pharmacodynamics, there are no recommended weak opioid agents for the treatment of neuropathic pain.

**Acetaminophen (Tylenol)** 500-1000 mg PO q6-8 hours (max 3 grams/24hrs)

### Is pain now adequately controlled?

Yes: Reassess at least monthly using the [Follow-Up Pain Assessment Tool](#)

Inadequate: ADD Strong opioid (Step 3)

## ► Step 3: Strong Opioid

Consider completing an opioid risk tool and order a **bowel routine** (See: [Constipation Guideline](#)). Start with low doses and titrate slowly to effect.

**Hydromorphone (Dilaudid)** 0.5 - 1.0mg PO q4h (or 0.2 mg SC). Due to the accumulation of metabolites, monitor closely for toxicity.

**Fentanyl Transdermal Patch** (for controlled pain) 12 mcg/h q72hours. Not recommended in opioid naïve patients ([Opioid Conversion Table](#)). Also available SL/SC.

**Buprenorphine Transdermal Patch** (for controlled pain) 5 mcg/h q7days ([Opioid Conversion Table](#)). Is not covered by Alberta Blue Cross. Access may be limited.

**Methadone** 1-2 mg/day PO (consider referral to Palliative Care; requires specific education and licensing by CPSA)

### Is pain now adequately controlled?

Yes: Reassess at least monthly using the [Follow-Up Pain Assessment Tool](#)

No: [Refer to Palliative Care](#)

## Considerations for Opioid Titration:

- Ongoing pain re-assessment is **critical**.
- Titrate analgesics every 3-7 days as needed and tolerated. Slower titration may be required.
- **Titration up the regular Opioid dose:**
  1. Add the total amount of opioid used in the last 24 hours (regular and breakthrough doses). Divide the total dose by 6, and prescribe this amount every 4 hours (q4H).

**OR**

  2. For ongoing pain exceeding patients pain control targets, adjust as follows:
    - For pain rated 3-6, increase dose of opioid by 25%
    - For pain rated 7-10, increase dose of opioid by 50%
- **Breakthrough (PRN) dose prescription:** 10% of total 24 hour opioid dose q 1-2 hrs PRN.
- If the patient is also taking benzodiazepines, consider titrating down the dose, while opioids are being increased. If not, titrate opioids more slowly.

**Considerations at End of Life:** In the last days of life, see the [End of Life Pain Algorithm](#)

## Conservative Kidney Management Acronym Legend

Acronym:	Intended Meaning:
ATC	Around the Clock
BID	Twice Daily
CKD	Chronic Kidney Disease
CKM	Conservative Kidney Management
COPD	Chronic Obstructive Pulmonary Disease
CO <sub>2</sub>	Carbon Dioxide
EOL	End of Life
ESA	Erythropoietin Stimulating Agent
ESKD	End Stage Kidney Disease
GFR	Glomerular Filtration Rate
GI	Gastrointestinal
g/L	Grams per litre
HgB	Hemoglobin
IN	Intranasal
IU	International Units
IV	Intravenous
kg	Kilogram
mcg	Microgram
mg	Milligram
mL	Millilitre

Acronym:	Intended Meaning:
mmol/L	Millimoles per Litre
OTC	Over the Counter
PO	By Mouth
PRN	As Needed
NSAID	Non-steroidal Anti-inflammatory Drugs
q(1-8)d	Every (Time Eg, 2) Days
q(1-8)h	Every (Time Eg, 4) Hours
q(1-8)weeks	Every (Time Eg. 2) Weeks
QHS	At Bedtime
RLS	Restless Leg Syndrome
SC	Subcutaneous
SL	Sublingual
SNRI	Serotonin and Norepinephrine Reuptake Inhibitors
SSRI	Selective Serotonin Reuptake Inhibitors
TCA	Tricyclic Antidepressant
TID	Three Times a Day
>	Greater Than
≥	Greater Than or Equal To
<	Less Than
≤	Less Than or Equal To